arXiv:astro-ph/9711044v1 6 Nov 1997

Keck Spectroscopy of Three Gravitational Lens Systems Discovered in the JVAS and CLASS Surveys

Christopher D. Fassnacht & Judith G. Cohen
Palomar Observatory, California Institute of Technology, 105-24, Pasadena, CA 91125
cdf@astro.caltech.edu, jlc@astro.caltech.edu

ABSTRACT

We present spectra of three gravitational lens systems taken with the Low Resolution Imaging Spectrograph on the W. M. Keck Telescopes. All of the systems were discovered in the JVAS and CLASS radio surveys, which were designed to find lenses suitable for measuring H_0. Previous spectra of these systems had low signal-to-noise ratios, and only one of the source redshifts was secure. Our observations give unambiguous lens and source redshifts for all of the systems, with $(z_\ell,z_s) = (0.4060,1.339)$, $(0.5990,1.535)$ and $(0.4144,1.589)$ for B0712+472, B1030+074 and B1600+434, respectively. The observed image splittings in the systems imply that the masses of the lensing galaxies within their Einstein rings are 5.4×10^{10}, 1.2×10^{11}, and $6.3\times10^{10} h^{-1} M_\odot$. The resulting V-band mass-to-light ratios for B0712+472 and B1030+074, measured inside their Einstein ring radii, are $\sim 10h\ (M/L)_{\odot,V}$, slightly higher than values observed in nearby ellipticals. For B1600+434, the mass-to-light ratio is $48h\ (M/L)_{\odot,V}$. This high value can be explained, at least in part, by the prominent dust lane running through the galaxy. Two of the three lens systems show evidence of variability, so monitoring may yield a time delay and thus a measurement of H_0.

Subject headings: distance scale — galaxies: distances and redshifts — gravitational lensing — quasars: individual (B0712+472, B1030+074, B1600+434)

1. Introduction

It has been known for many years that gravitational lens systems can in principle be used to determine the values of various cosmological parameters. Refsdal (1964) described a method by which variations in intensity of a multiply-imaged background source could be used to determine the Hubble Constant. A well-constrained model of the lensing potential can be used to predict differences in light travel time along the multiple paths from source to observer; knowing the redshifts of the background source and the lensing galaxy for each system is crucial to this method. The ratio of the observed to the predicted time delays is directly proportional to H_0. Time delays have been measured for two lens systems to date: 0957+561 (Kundić et al. 1995; Kundić et al. 1997a; Oscoz et al. 1997) and 1115+080 (Schechter et al. 1997; Bar-Kana 1997). Combining the measured time delays with current models for these systems has given values of H_0 in the range ≈ 60–70 km/sec/Mpc for 0957+561 (Kundić et al. 1997a; Oscoz et al. 1997; Falco et. al 1997) and ≈ 40–55 km/sec/Mpc for 1115+080 (Schechter et al. 1997; Keeton & Kochanek 1997; Kundić et al. 1997b; Courbin et al. 1997). Improvements in the time delay measurements and additional constraints on the lens models are expected. The value of Refsdal's method will be proved if several lenses can be used to measure H_0 and give consistent values.

The Jodrell-VLA Astrometric Survey (JVAS; Patnaik et al. 1992; Browne et al. 1997b) and the Cosmic Lens All-Sky Survey (CLASS; Jackson et al. 1995; Myers et al. 1995; Browne et al. 1997a; Myers et al. 1997) contain observations of ∼10,000 flat-spectrum radio sources. One of the primary goals of these surveys is to find gravitational lens systems which can be used to measure H_0. To date, 11 new gravitational lenses have been discovered in the JVAS and CLASS surveys, and 16 lens candidates are being investigated further. The lens and source redshifts have been determined for almost all of the newly discovered lens systems. This paper presents spectra taken at the W. M. Keck Observatory of three of the systems with missing redshifts, B0712+472, B1030+074 and B1600+434. There is evidence of variability in B0712+472 (Jackson et al. 1997b) and B1600+434 (Jaunsen & Hjorth 1997), so at least two of the systems present the possibility of being used to measure H_0.

We use $H_0 = 100\ h$ km s^{-1} Mpc^{-1} and assume $q_0 = 0.5$ throughout this paper.

2. Targets

Below we present information on previous observations of the lens systems that are discussed in this paper.

2.1. B0712+472

This lens system consists of four images of the background source in a typical lensing geometry (e.g. Blandford & Narayan 1992). The maximum image separation is $1\farcs27$. Radio maps made with the VLA and MERLIN do not resolve the four images (Jackson et al. 1997b). Images taken with WFPC2 on HST show all four images and, in addition, the lensing galaxy (Jackson et al. 1997a). Photometry derived from the WFPC2 images give the lens magnitude as $V \sim 20.2$ and $I \sim 20.0$[1], measured in an elliptical aperture with major and minor axes of 2 and $1''$, respectively (Jackson et al. 1997b). The total source magnitudes, assuming the images were point sources, were $V \sim 23$ and $I \sim 22.5$. Low signal-to-noise ratio optical spectra of the system, taken with the William Herschel Telescope (WHT), show weak broad emission lines (C IV, C III], Mg II) giving a source redshift of $z_s \sim 1.33$, and a "hint of Ca H and K" absorption, which tentatively gives $z_\ell \sim 0.40$ (Jackson et al. 1997b). The redshifts derived from the WHT spectra, especially that of the lens, are not secure. However, both are confirmed to be correct by the observations described in this paper.

2.2. B1030+074

The B1030+074 system consists of two flat-spectrum radio components separated by $1\farcs56$ with a flux ratio of 15 to 1 (Browne et al. 1997a; Xanthopoulos et al. 1997). These components are unresolved at the milliarcsecond resolution of VLBI. The system was imaged with WFPC2 and both images of the background source are seen, as is the lensing galaxy. Component A, the brighter image, has magnitudes of $V \sim 20$ and $I \sim 19$ and component B is ∼2.5 − 3 magnitudes fainter in each band; the lensing galaxy is estimated to have magnitudes $I \sim 20.5$ and $V \sim 22$

---

[1] In this paper all WFPC2 F555W and F814W magnitudes have been converted to Johnson V and Cousins I magnitudes, respectively, using transformations in Holtzman et al. (1995).

(Xanthopoulos et al. 1997). Spectra taken with the COSMIC spectrograph (Kells et al. 1997) on the Hale Telescope show no clear spectral features (Vermeulen & Womble, private communication).

2.3. B1600+434

This system is also a double, with separation 1.″4 and a flux ratio of 1.30 to 1 at radio wavelengths (Jackson et al. 1995). Jaunsen & Hjorth (1997) have imaged the system with the Nordic Optical Telescope and find that the lensing galaxy is an edge-on spiral with $B = 23.6, V = 22.0, R = 21.1$, and $I = 20.3$. The background source images have total magnitudes of $B = 21.9, V = 21.7, R = 21.1$, and $I = 20.4$. Imaging with WFPC2 has shown that the lensing galaxy has a prominent dust lane along its major axis (Jackson et al. 1997a). Previous spectroscopy with the WHT detected broad C IV, C III] and Mg II emission lines from the background source. These lines give a source redshift of $z_s = 1.61$, but no redshift for the lens was determined (Jackson et al. 1995). Jaunsen & Hjorth (1997) used the lensing galaxy colors to get a photometric redshift of $z \sim 0.4$, which is confirmed by the observations presented in this paper.

3. Observations and Data Reduction

All three of the systems were observed with the Low Resolution Imaging Spectrograph (LRIS; Oke et al. 1995) in longslit mode on the Keck Telescopes. The 300 g/mm grating was used, giving a scale of 2.44 Å/pix. The longslit was aligned with the major axis of the lensing galaxy (B0712+472 and B1600+434) or the axis defined by the two images of the background source (B1030+074). All objects were observed for 3000 sec total integration time; other details of the observations are given in Table 1. The data were reduced using standard IRAF[2] routines. The bias levels were estimated using the overscan region on the chip. For the observations of B1600+434, the flat-field frame was constructed from dome flats; in all other cases, exposures with the internal flat-field lamp were used. The spectra were extracted using the IRAF implementation of the "optimal extraction" technique described in Horne (1986) and Marsh (1989). For the 1997 Feb 07 observation of B0712+472 and the 1997 Feb 14 observation of B1030+074, the emission from the background source and the lensing galaxy were spatially separated on the slit, so two spectra were extracted for each system. Wavelength calibration was performed using sky lines (1997 Feb 07) or arc lamps taken after each science exposure (all other observations). Observations of the Oke spectrophotometric standard stars G191B2B and HZ44 (Oke 1990) were used to remove the response function of the chip. For the data taken on 1996 June 18 and 1997 Feb 14, exposures of BL Lac objects were used to remove atmospheric absorption features. The individual spectra for each object were weighted by their signal-to-noise ratios and combined.

4. Results

The final spectra are shown in Figures 1 – 5. For each lens system, the spectral features (Table 2) can be explained as the sum of a lens at z_ℓ and a background source at z_s, where a single choice of z_ℓ and z_s (Table 3) suffices. Uncertainties in the redshifts were estimated by taking the RMS scatter in the redshifts calculated from the individual spectral lines. For B1600+434 the source redshift uncertainty was estimated as half the difference of the redshifts calculated from the two broad emission lines. Further discussion of the individual systems follows below.

4.1. B0712+472

The B0712+472 spectra show a lensing galaxy at $z_\ell = 0.4060$ and a background source at $z_s = 1.339$. These results confirm the Jackson et al. (1997b) tenta-



Fig. 1.— Spectrum of lensing galaxy in B0712+472 system taken on 1997 Feb 06.

---

[2] IRAF (Image Reduction and Analysis Facility) is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy (AURA) under cooperative agreement with the NSF.

TABLE 1
OBSERVATIONS

Source	Date	Telescope	t_{exp} (sec)	Slit Width ($''$)	P.A.	Coverage (Å)
B0712+472	1997 Feb 06	Keck II	3000	0.7	75	3593–8588
	1997 Feb 14	Keck II	3000	0.7	79	4208–9206
B1030+074	1997 Feb 14	Keck II	3000	1.0	322	4211–9208
B1600+434	1996 Jun 18	Keck I	3000	1.0	−120	4606–9358



Fig. 2.— Spectrum of background source in B0712+472 system taken on 1997 Feb 06. The spectrum also includes a significant amount of light from the extended lensing galaxy. The C III], C II] and Mg II are associated with the background source. All other lines are associated with the lensing galaxy.



Fig. 3.— Spectrum of lensing galaxy in B1030+074 system taken on 1997 Feb 14.



Fig. 4.— Spectrum of background source in B1030+074 system taken on 1997 Feb 14. The C III], C II] and Mg II emission lines are associated with the background source. All other lines are associated with the lensing galaxy.



Fig. 5.— Spectrum of B1600+434 containing light from both the lens and the source. The C III] and Mg II emission lines are associated with the background source. All other lines are associated with the lensing galaxy.

tive redshifts. The lensing galaxy has a typical early-type galaxy spectrum (e.g. Kennicutt 1992). The moderately strong 4000Å break, small equivalent-width Balmer absorption lines and lack of O II emission indicate that little star formation is occuring in this galaxy. The background source shows broad C III] and Mg II emission lines typical of a quasar spectrum.

4.2. B1030+074

The B1030+074 system contains a lensing object at $z_\ell = 0.5990$ and a background source at $z_s = 1.535$. In this system as well, the lensing object spectrum is typical of an early-type galaxy. The background source shows broad C III] and Mg II emission lines. The broad emission lines have low equivalent widths, raising the possibility that the background source is a BL Lac-type object. It should be noted, however, that the source spectrum is almost certainly contaminated by some light from the lensing object, which will raise the continuum level and reduce the apparent equivalent widths of the lines.

4.3. B1600+434

The B1600+434 spectrum has the lowest signal-to-noise ratio of the observations presented in this paper. However, both lens and source redshifts ($z_\ell = 0.4144, z_s = 1.589$) are clearly determined. The background source is a quasar with broad C III] and Mg II emission lines, as was seen in Jackson et al. (1995). The lens spectrum is indicative of a later galaxy type than that observed in the B0712+472 and B1030+074 systems. This spectrum has [O II] emission and a less prominent 4000Å break. This is consistent with images of the lensing galaxy which show that it is an edge-on spiral (Jaunsen & Hjorth 1997; Jackson et al. 1997a). For later calculations, we will assume that the lensing object is an Sa or Sb galaxy. However, determining the galaxy type from the spectrum may be complicated by extinction from the dust lane, which runs the length of the observable disk (Jackson et al. 1997a).

5. Discussion

In order to use a lens system to measure H_0, it is necessary to first know the source and lens redshifts. These redshifts are used to derive the angular diameter distances to the source and the lens, which have an inverse dependence on H_0. The observed image positions and other information can then be used to construct a model of the lensing potential. The lens model is combined with the angular diameter distances to give predicted time delays:

$$\Delta t_i = (1+z_\ell)\frac{D_\ell D_S}{cD_{\ell s}}\left[\frac{1}{2}|\theta_i - \beta|^2 - \phi(\theta)\right],$$

where D_ℓ, D_S and $D_{\ell s}$ are the angular diameter distances to the lens, to the source, and between the lens and source, respectively; θ_i is the image position; β is the source position and ϕ is the scaled lensing potential (e.g. Hogg & Blandford 1994). The predicted time delay is proportional to h^{-1} from the ratio of angular diameter distances. Thus, if the background source is variable and time delays can be measured, the ratio between the observed and predicted delays will give h.

We can also use these lenses to study the properties of galaxies at moderate redshifts. For example, the extent of the image-splitting by the lens gives a dircect estimate of the mass inside the Einstein ring of the lens. For the lens systems discussed in this paper, the Einstein ring radii correspond to physical sizes of $2-3h^{-1}$ kpc. The mass is estimated as (e.g. Blandford & Narayan 1992):

$$M_E \sim 1\times 10^{12}\left(\frac{D}{1\ \mathrm{Gpc}}\right)\left(\frac{\theta_E}{3\ ''}\right)^2 M_\odot, \quad D \equiv \frac{D_\ell D_s}{D_{\ell s}},$$

where θ_E is the angular radius of the Einstein ring. For these lenses, we find values of $M_E \sim 5\times 10^{10} - 1\times 10^{11} h^{-1} M_\odot$.

With the above masses and the photometry for these systems, we can compute approximate mass-to-light ratios for the lensing galaxies. For B0712+472 and B1030+074, the lens magnitudes discussed in Section 2 were taken within apertures roughly the size of the Einstein rings in the systems, so we use those values in the following calculations. For B1600+434 we have used the WFPC2 images to estimate the lens magnitude in the appropriate aperture, finding $I \sim 21.8$. We converted the observed I-band magnitudes to rest-frame V-band magnitudes using k-corrections for the lens redshifts and rest-frame $(V-I)$ colors for typical E/S0 (for B0712+472 and B1030+074) and Sa (B1600+434) galaxies. The resulting mass-to-light ratios for B0712+472 and B1030+074 are $\sim 10h(M/L)_{\odot,V}$. For B1600+434 we find $(M/L)_E = 48h(M/L)_{\odot,V}$, confirming the value found by Jaunsen & Hjorth (1997).

The values of mass-to-light for B0712+472 and B1030+074 are slightly higher than mass-to-light ratios of nearby ellipticals within the same physical radii (e.g. van der Marel 1991; Gerhard et al. 1997). We note that we are biased toward finding high mass systems when looking for gravitational lenses, since these systems have a larger cross-section for lensing. Also, the slightly different apertures used to compute the masses and luminosities may be biasing the results toward higher mass-to-light ratios. In addition, the presence of dust in the lensing galaxies would cause some of the light to be lost, thereby increasing the observed mass-to-light ratio. The dust lane in B1600+434 is clearly seen and is responsible, at least in part, for the very high mass-to-light ratio calculated for this system. There is also evidence for absorption by dust in B0712+472 (Jackson et al. 1997a). Evidence for dust has been seen in other lens systems as well (Larkin et al. 1994; Malhotra et al. 1997), indicating that dusty lenses may be quite common. Infrared imaging of these systems would provide more accurate measurements of the mass-to-light ratios.

6. Prospects for Measuring H_0

In order to use these systems to measure H_0, well-constrained models of the lensing potentials must be constructed. The following observations could provide data that would aid in the modelling: (1) detecting milliarcsecond-scale structure in the radio images, (2) determining the velocity dispersion of the lensing galaxy or, in the case of B1600+434, measuring the galaxy rotation curve, and (3) imaging the systems at high-resolution with NICMOS to provide accurate positions of the lensed images with respect to the lensing galaxy, unbiased by dust extinction.

The final piece of the H_0 puzzle is measuring time delays between the lensed images. Jackson et al. (1997b) find slight variability in the component fluxes in B0712+472, with a ∼30% overall decrease in flux between 1995 and 1996. The background source in B1600+434 appears to be variable as well. Our VLA observations of B1600+434 show that between 1995 and 1996, the flux densities of the two components decreased by ∼50%, and the component flux ratio changed from 1.30 to 1.11. In addition, Jaunsen and Hjorth (1997) see optical variations in B1600+434. Thus, at least two of the sources show variability and, if observed regularly, present the possibility of being used to measure H_0.

We thank Nicole Vogt, Tony Readhead and the anonymous referee for insightful comments on the manuscript, and are grateful to Lori Lubin, Mike Pahre and David Hogg for many helpful discussions. We are indebted to Terry Stickel, Wayne Wack, Chuck Sorenson and the Keck staff for assistance with the observations. The W. M. Keck Observatory was made possible by a generous grant from the W. M. Keck Foundation. This work is supported by the NSF under grant #AST 9420018.

REFERENCES

Bar-Kana, R. 1997, astro-ph/9701068

Blandford, R. D. & Narayan, R., 1992, ARA&A, 30, 311

Browne, I. W. A. et al. 1997, in Observational Cosmology with the New Radio Surveys, eds., M. Bremer, N. Jackson & I. Perez-Fournon, (Dordrecht: Kluwer) 305

Browne, I. W. A., Patnaik, A. R., Wilkinson, P. N. & Wrobel, J. M. 1997, MNRAS, in press

Courbin, F., Magain, P., Keeton, C. R., Kochanek, C. S., Vanderriest, C., Jaunsen, A. O., & Hjorth, J. 1997, A&A, 342, L1

Falco, E. E., Shapiro, I. I., Moustakas, L. A., & Davis, M. 1997, ApJ, 484, 70

Gerhard, O., Jeske, G., Saglia, R. P., Bender, R. 1997, MNRAS, astro-ph/9710129

Hogg, D. W. & Blandford, R. D. 1994, MNRAS, 268, 889

Holtzman, J. A., Burrows, C. J., Casertano, S., Hester, J. J., Trauger, J. T., Watson, A. M., & Worthey, G. 1995, PASP, 107, 1065

Horne, K. 1986, PASP, 98, 609

Jackson, N., et al. 1995, MNRAS, 274, L25

Jackson, N. J., Nair, S., Browne, I. W. A. 1997, in Observational Cosmology with the New Radio Surveys, eds., M. Bremer, N. Jackson & I. Perez-Fournon, (Dordrecht: Kluwer) 315

Jackson, N. J. et al. 1997, MNRAS, in press

Jaunsen, A. O. & Hjorth, J. 1997, A&A, 317, L39

TABLE 2
DETECTED SPECTRAL LINES

			Observed Wavelength		
Ion	λ_0 (Å)	Object	B0712+472	B1030+074	B1600+434
C III]	1909	Source	4461	4834	4930
C II]	2326	Source	5448	5898	···
Mg II	2796	Source	6540	7095	7252
Mg II	2796	Lens	···	4469	···
Mg II	2802	Lens	···	4480	···
Mg I	2852	Lens	···	4563	···
O II	3727	Lens	···	5963	5274
Hη	3835	Lens	5391	6134	5418
Hζ	3889	Lens	···	6212	5498
Ca II K	3934	Lens	5529	6288	5566
Ca II H[a]	3968	Lens	5579	6345	5615
Hδ	4102	Lens	5768	6560	5802
G-band	4300	Lens	6053	6887	6094
Hγ	4340	Lens	6104	6943	6143
Hβ	4861	Lens	6839	···	6876
Mg I	5172	Lens	7266	···	···
Mg I	5183	Lens	7287	···	···
Fe I	5269	Lens	7409	···	···
Ca I	5589	Lens	7863	···	···
Na I D	5890	Lens	8283	···	···
Na I D	5896	Lens	8290	···	···

[a] May be blended with Hϵ λ3970.

TABLE 3
LENS SYSTEM PARAMETERS

System	z_ℓ	z_s	D_ℓ (h^{-1} Mpc)	D_s (h^{-1} Mpc)	$D_{\ell s}$ (h^{-1} Mpc)	M_E[a] (h^{-1} $M_\odot$)	$(M/L)_V$[b] (h $(M/L)_\odot$)
B0712+472	0.4060 ± 0.0002	1.339 ± 0.002	668 ± 0.16	887 ± 0.042	486 ± 0.34	$5.4 \pm 0.22 \times 10^{10}$	8.6 ± 0.93
B1030+074	0.5990 ± 0.0003	1.535 ± 0.003	784 ± 0.13	879 ± 0.16	385 ± 0.46	$1.2 \pm 0.050 \times 10^{11}$	11 ± 1.2
B1600+434	0.4144 ± 0.0003	1.589 ± 0.006	675 ± 0.24	876 ± 0.36	507 ± 0.53	$6.3 \pm 0.25 \times 10^{10}$	48 ± 5.2

[a] Uncertainties calculated assuming a 1% uncertainty in image separation.

[b] Uncertainties calculated assuming a 0.1 magnitude uncertainty in photometry.

Keeton, C. R. & Kochanek, C. S. 1997, ApJ, 487, 48

Kells, W. P., Sivaramakrishnan, A., Dressler, A., Epps, H., Hilyard, G, & Pardillian, G. 1997, in preparation

Kennicutt, R. C., Jr. 1992, ApJS, 79, 255

Kundić, T. et al. 1995, ApJ, 455, L5

Kundić, T. et al. 1997a, ApJ, 482, 75

Kundić, T., Cohen, J. G., Blandford, R. D., & Lubin, L. M. 1997b, AJ, 114, 507

Larkin, J. E, et al. 1994, ApJ, 420, L9

Malhotra, S., Rhoads, J. E., Turner, E. L. 1997, MNRAS, 228, 138

Mao & Schneider 1997, astro-ph/9707187

Marsh, T. 1989, PASP, 101, 1032

Myers, S. T., et al. 1995, ApJ, 447, L5

Myers, S. T., et al. 1997, in preparation

Oke, J. B. 1990, AJ, 99, 1621

Oke, J. B. et al. 1995, PASP, 107, 375

Oscoz, A., Mediavilla, E., Goicoechea, L. J., Serra-Ricart, M. & Buitrago, J. 1997, ApJ, 479, L89

Patnaik, A. R., Browne, I. W. A., Wilkinson, P. N., & Wrobel, J. M. 1992, MNRAS, 254, 655

Refsdal, S. 1964, MNRAS, 128, 307

Schechter, P. L, et al. 1997, ApJ, 475, L85

van der Marel, R. P. 1991, MNRAS, 253, 710

Xanthopoulos, E., et al. 1997, in preparation

This 2-column preprint was prepared with the AAS LaTeX macros v4.0.